



PCT/EP200 4 / 0 0 4 1 2 2



16.04.04

INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 11 MAY 2004


WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *Andrew*

Dated 24 February 2004

Patents Form 1/77

Patents Act 1977
(Rule 16)

The
Patent
Office



PCT/EP 2004/009122

16.04.04

1/77

Request for grant of a patent

(See the notes on the back of this form. You can also get a free explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your Reference **AP/PB60212P**

19APR03 EB01248-1 D01030
P01/7700 0.00-0308968.7

2. Patent application number
(The Patent office will fill in this part)

3. Full name, address and postcode of the or of each applicant (underline all surnames)

**GLAXO GROUP LIMITED
GLAXO WELLCOME HOUSE
BERKELEY AVENUE
GREENFORD
MIDDLESEX
UB6 0NN
GB**

0308968.7

00473587003

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its corporation

GB

4 Title of the invention **MEDICAMENTS**

5 Name of your agent (if you know one) **PETER I DOLTON**

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

**GLAXOSMITHKLINE
CORPORATE INTELLECTUAL PROPERTY
CN925.1
980 GREAT WEST ROAD
BRENTFORD
MIDDLESEX
TW8 9GS, GB**

Patents ADP number (if you know it)

08321846001

6. If you are declaring priority from one or more earlier patent applications, give the country and date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country	Priority application number (if you know it)	Date of Filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant a patent required in support of this request? (Answer yes if:

a) any applicant named in part 3 is not an inventor, or

b) there is an inventor who is not named as an applicant, or

c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document.

Continuation sheets of this form

Description 18

Claim(s) 1

Abstract 1

Drawing(s) -

10. If you are also filing any of the following, state how many against each item

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patent Form 9/77)

Request for substantive examination (Patent Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application

P. I. Donon

Signature PETER I DONON 17 April 2003
AGENT FOR THE APPLICANTS

12. Name and daytime telephone number of person to contact in the United Kingdom

JEAN HARNEY
020 8047 4420

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication of communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the patent Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been received

a) Notes

If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.

b) Write your answers in capital letters using black ink or you may type them.

c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form

If you have answered "Yes" Patents Form 7/77 will need to be filed.

d) Once you have filled in the form you must remember to sign and date it.

e) For details of the fee and ways to pay please contact the Patent Office.

MEDICAMENTS

The present invention relates to therapeutic combinations comprising paroxetine and a NK₁ receptor antagonist, to pharmaceutical compositions containing said combinations and their use in the treatment of depression and /or anxiety.

Paroxetine ((-)-trans-4-(4'-fluorophenyl)-3-(3'-4'-methylenedioxyphenoxymethyl) piperidine) and its salts are commercially available and approved for use in humans for treatment and prophylaxis of, *inter alia*, anxiety, depression, obsessive compulsive disorder (OCD) and panic.

NK₁ receptor antagonists are known to be useful in the treatment of anxiety and depression, chemotherapy-induced nausea and vomiting and post-operative nausea and vomiting. Preclinical data suggest that NK₁ receptor antagonists may be useful in a variety of other disorders including pain, inflammatory diseases, allergic disorders, CNS disorders, skin disorders, cough and gastrointestinal disorders.

US 6117855 describes the use of a CNS-penetrant NK₁ receptor antagonist and an antidepressant or anti-anxiety for the manufacture of a medicament for the treatment or prevention of depression and/or anxiety.

There is however no specific disclosure of such combinations with paroxetine.

WO 01/25219 and WO 02/32867 teach that the NK₁ receptor antagonists described therein may be administered in combination with a SSRI agent. However, there is no specific teaching concerning the use of a combination of paroxetine with the compounds claimed therein in the treatment of depression and /or anxiety.

We have found that therapeutic compositions comprising a combination of paroxetine and a NK₁ receptor antagonist provide a useful and unexpectedly advantageous combination for the treatment of depression and/or anxiety.

In particular, it has now been found that by combining paroxetine and a NK₁ receptor antagonist, a synergistic anti-depressive and/or anti-anxiety effect is achieved.

Furthermore, the synergistic effect is also observed when each component is employed in the combination in an amount or dosage which is less than that normally expected to produce a fully effective therapeutic response when the component is administered alone.

- 5 It is a feature of this invention that the use of such a combination will provide one or more of the following effects: the synergistic effect of the combination providing a more efficacious anti-depressive and/or anti-anxiety drug and/or a better tolerated drug treatment and/or a more rapid onset of the anti-depressive and/or anti-anxiety activity.
- 10 When used in any of the contexts or aspects of the present invention, paroxetine may be administered as the free base, or in the form of any physiologically acceptable salt thereof, including all hydrated or anhydrous forms and all polymorphic forms of such salts. In particular, references to paroxetine include, without limitation, paroxetine hydrochloride, paroxetine hydrochloride hemihydrate, paroxetine hydrochloride anhydrate, paroxetine
- 15 mesylate and all polymorphic forms thereof.

According to one aspect of the invention, there is provided a combination comprising paroxetine and a NK₁ receptor antagonist.

- 20 NK₁ receptor antagonists for use in the present invention include those generically and specifically disclosed in the following patent specifications whose disclosures are incorporated by reference;

European Patent Specification Nos. 284942, 327009, 333174, 336230, 360390, 394989, 428434, 429366, 436334, 443132, 446706, 482539, 484719, 499313, 512901, 512902, 25 514273, 514275, 517589, 520555, 522808, 525360, 528495, 532456, 533280, 577394, 591040, 615751, 684257, 1176144, 1110958, 1176144, 1172106, 1103545, and 1256578; and in

International Patent Specification Nos. 90/05525, 90/05729, 91/02745, 91/12266, 30 91/18016, 91/18899, 92/01688, 92/06079, 92/15585, 92/17449, 92/20676, 92/21677, 92/22569, 93/00331, 93/01159, 93/01160, 93/01165, 93/01169, 93/01170, 94/01402, 94/26735, 95/06645, 95/08549, 95/14017, 95/16679, 95/18124, 95/23798, 95/28389, 95/33744, 96/05181, 96/18643, 96/21661, 96/29326, 96/32386, 96/34857, 96/37489, 97/02824, 97/05110, 97/08166, 97/13514, 97/14671, 97/16440, 97/17362, 97/19074, 35 97/19084, 97/19942, 97/21702, 97/22597, 97/22604, 97/23455, 97/24324, 97/24350, 97/25322, 97/25988, 97/27185, 97/30989, 97/30990, 97/30991, 97/32865, 97/38692,


97/44035, 97/49393, 97/49710, 98/02158, 98/04561, 98/07694, 98/07722, 98/08826,
98/13369, 98/17276, 98/18761, 98/18785, 98/18788, 98/20010, 98/24438, 98/24439,
98/24440, 98/24441, 98/24442, 98/24442, 98/24443, 98/24444, 98/24445, 98/24446,
98/24447, 98/28297, 98/43639, 98/45262, 98/49170, 98/54187, 98/57954, 98/57972,
5 99/00388, 99/01444, 99/01451, 99/07677, 99/07681, 99/09987, 99/21823, 99/24423,
99/25364, 99/26924, 99/27938, 99/36424, 99/52903, 99/59583, 99/59972, 99/62893,
99/62900, 99/64000, 00/02859, 00/06544, 00/06571, 00/06572, 00/06578, 00/06580,
00/15621, 00/20003, 00/21512, 00/21564, 00/23061, 00/23062, 00/23066, 00/23072,
00/20389, 00/25745, 00/26214, 00/26215, 00/34243, 00/34274, 00/39114, 00/47562,
10 01/77069, 01/25233, 01/30348, 01/87866, 01/94346, 01/90083, 01/87838, 01/85732,
01/77100, 01/77089, 01/77069, 01/46176, 01/46167, 01/44200, 01/32625, 01/29027,
01/25219, 02/32865, 02/00631, 02/81461, 02/92604, 02/38575, 02/57250, 02/22574,
02/74771, 02/26710, 02/28853, 02/102372, 02/85458, 02/81457, 02/74771, 02/62784,
02/60898, 02/60875, 02/51848, 02/51807, 02/42280, 02/34699, 02/32867, 02/32866,
15 02/26724, 02/24673, 02/24629, 02/18346, 02/16344, 02/16343, 02/16324, 02/12168,
02/08232 and 02/06236; and in

US Patent Specification Nos. 4839465, 5338845, 5594022, 6169097, 6197772, 6222038,
6204265, 6329392, 6316445, 2001039286, 2001034343, 2001029297, 2002193402,
20 2002147212, 2002147207, 2002143003 and 2002022624; and in

British Patent Specification Nos. 2216529, 2266529, 2268931, 2269170, 2269590,
2271774, 2292144, 2293168, 2293169 and 2302689; and in

25 Japanese Patent Specification No 6040995.

A particularly useful class of NK1 receptor antagonists for use in the combinations of the
present invention is represented by those compounds described in WO 0125219 having
the general formula(I)



wherein

R represents a halogen atom or a C₁₋₄ alkyl group;

R₁ represents hydrogen or a C₁₋₄ alkyl group;

- 5 R₂ represents hydrogen, a C₁₋₄ alkyl, C₂₋₆ alkenyl or a C₃₋₇ cycloalkyl group; or R₁ and R₂ together with nitrogen and carbon atom to which they are attached respectively represent a 5 to 6 membered heterocyclic group;

R₃ represents a trifluoromethyl, a C₁₋₄ alkyl, a C₁₋₄ alkoxy, a trifluoromethoxy or a halogen group;

- 10 R₄ represents hydrogen, a (CH₂)_qR₇ or a (CH₂)_rCO(CH₂)_pR₇ group;

R₅ represents hydrogen, a C₁₋₄ alkyl or a COR₆ group;

R₆ represents hydrogen, hydroxy, amino, methylamino, dimethylamino a 5 membered heteroaryl group containing 1 to 3 heteroatoms selected from oxygen, sulphur and nitrogen or a 6 membered heteroaryl group containing 1 to 3 nitrogen atoms;

- 15 R₇ represents hydrogen, hydroxy or NR₈R₉ wherein R₈ and R₉ represent independently hydrogen or C₁₋₄ alkyl optionally substituted by hydroxy or by amino;

R₁₀ represents hydrogen, a C₁₋₄ alkyl group or

R₁₀ together with R₂ represents a C₃₋₇ cycloalkyl group;

m is zero or an integer from 1 to 3;

- 20 n is zero or an integer from 1 to 3;

both p and r are independently zero or an integer from 1 to 4;

q is an integer from 1 to 4;

provided that, when R₁ and R₂ together with nitrogen and carbon atom to which they are attached respectively represent a 5 to 6 membered heterocyclic group:

- 25 i) m is 1 or 2;

ii) when m is 1, R is not fluorine and


iii) when m is 2, the two substituents R are not both fluorine and pharmaceutically acceptable salts and solvates thereof.

Within this class, particularly preferred compounds include those wherein R is selected independently from halogen or methyl, R₃ is trifluoromethyl both at the 3 and 5 position, R₁ is hydrogen or methyl, R₂ is hydrogen, methyl, 2-propenyl, or a cyclopropyl group or together with R₁ is a 3,6-dihydro-2H-pyridin-1-yl, a piperidin-1-yl or a pyrrolidin 1-yl group,
 5 R₁₀ represents hydrogen, a methyl or R₁₀ together with R₂ is a cyclopropyl group, R₄ is hydrogen, an aminoacetyl or amino ethyl group and R₅ is hydrogen or a methyl group.

A particularly preferred compound of formula (I) for use in the invention is

2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide and physiologically acceptable salts thereof
 10 (e.g. methansulphonate or hydrochloride).

Further examples of suitable NK1 receptor antagonists for use in the combinations of the present invention include those described in WO 0232867 having the general formula(II)



(II)

wherein

R represents a halogen atom or a C₁₋₄ alkyl group;

20 R₁ represents a C₁₋₄ alkyl group;

R₂ represents hydrogen or a C₁₋₄ alkyl group;

R₃ represents hydrogen, or a C₁₋₄ alkyl group;

R₄ represents a trifluoromethyl group;

R₅ represents hydrogen, a C₁₋₄ alkyl group or C(O)R₆;

25 R₆ represents C₁₋₄ alkyl, C₃₋₇ cycloalkyl, NH(C₁₋₄ alkyl) or N(C₁₋₄alkyl)₂;

m is zero or an integer from 1 to 3;


n is an integer from 1 to 3;

and pharmaceutically acceptable salts and solvates thereof.

A particularly preferred compound of formula (II) for use in the combinations of the invention is

- 4-(S)-(4-Acetyl-piperazin-1-yl)-2-(R)-(4-fluoro-2-methyl-phenyl)-piperidine-1-carboxylic acid, [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide and pharmaceutically acceptable salts (e.g. hydrochloride, methanesulphonate, sulphate, p-toluensulphonate) or solvates thereof.

Another further preferred class of Nk1 receptor antagonists for use in the present invention is that described in PCT /EP03/01308 having general formula(III)



wherein

R represents halogen or C₁₋₄ alkyl ;


R₁ represents C₁₋₄ alkyl;

- 15 R₂ or R₃ independently represent hydrogen or C₁₋₄ alkyl;

R₄ represents trifluoromethyl, C₁₋₄ alkyl, C₁₋₄ alkoxy, trifluoromethoxy or halogen;

R₅ represents hydrogen, C₁₋₄ alkyl or C₃₋₇ cycloalkyl;

R₆ is hydrogen and R₇ is a radical of formula (W):



(W)

- 20 or R₆ is a radical of formula (W) and R₇ is hydrogen;

X represents CH₂, NR₅ or O;

Y represents Nitrogen and Z is CH or Y represents CH and Z is Nitrogen;

A represents C(O) or S(O)_q, provided that when Y is nitrogen and Z is CH, A is not S(O)_q;

- 25 m is zero or an integer from 1 to 3;

n is an integer from 1 to 3;

p and q are independently an integer from 1 to 2;

and pharmaceutically acceptable salts and solvates thereof.

Particularly preferred compounds of formula (III) include those in which R₆ is hydrogen, R₇ is a radical of formula (W) and Y is CH and Z is nitrogen or wherein R₆ is a radical of formula (W), R₇ is a hydrogen and Y is nitrogen and Z is CH; A is C(O) and X is CH₂.

5

A particularly preferred compound of formula (III) for use in the invention is 2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide and pharmaceutically acceptable salts (e.g. hydrochloride, methanesulphonate or maleate) and solvates thereof.

10

The subject matter of the above identified patent applications is incorporated herein by reference.

15 A preferred embodiment of the invention provides a combination of paroxetine and a NK1 receptor antagonist selected from

2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide and physiologically acceptable salts thereof (e.g. methanesulphonate and hydrochloride);

20 2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide and physiologically acceptable salts thereof (e.g. hydrochloride, methanesulphonate or maleate);

25 2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide and physiologically acceptable salts thereof (e.g. hydrochloride, methanesulphonate or maleate);

2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl-benzyl)]-(2S-phenyl-piperidin-3S-yl)-amine.

30 A particularly preferred combination according to the invention comprises paroxetine and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide methanesulphonate.

35 A further particularly preferred combination according to the invention comprises paroxetine hydrochloride and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide methanesulphonate.

A further particularly preferred combination according to the invention comprises paroxetine hydrochloride hemihydrate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

A further particularly preferred combination according to the invention comprises paroxetine hydrochloride anhydrate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

A further particularly preferred combination according to the invention comprises paroxetine mesylate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

The present invention thus provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said animal with a therapeutically effective amount of a combination of paroxetine and a NK1 receptor antagonists.

A preferred embodiment of the invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said animal with a therapeutically effective amount of a combination of paroxetine and a NK1 receptor antagonists selected from

2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide and pharmaceutically acceptable salts thereof (e.g. methansulphonate and hydrochloride);

2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methylamide pharmaceutically acceptable salts (e.g. hydrochloride, methanesulphonate or maleate);

2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methylamide pharmaceutically acceptable salts (e.g. hydrochloride, methanesulphonate or maleate);

2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl-benzyl)]-(2S-phenyl-piperidin-3S-yl)-amine).

More particularly, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination of paroxetine and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide and physiologically acceptable salts thereof (e.g. methansulphonate and hydrochloride).

In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination of paroxetine and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination of paroxetine hydrochloride and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination of paroxetine hydrochloride hemihydrate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination of paroxetine hydrochloride anhydrate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

In a further preferred aspect, the present invention provides a method for the treatment of depression and/or anxiety in a mammal including a human, which comprises treating said mammal with a therapeutically effective amount of a combination of paroxetine mesylate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.

Reference herein to treatment extends to prophylaxis as well as the treatment of established depression and/or anxiety or symptoms.

5 As used herein, the term depression includes depressive disorders. Thus, for example, depressive disorders include Major Depressive Disorders (MDD), including bipolar depression, unipolar depression, single or recurrent major depressive episodes, recurrent brief depression, with or without psychotic features, catatonic features, melancholic features including anorexia, weight loss, atypical features, anxious depression,
10 cyclothymic or postpartum onset.

Other mood disorders encompassed within the term major depressive disorders include dysthymic disorders with early or late onset and with or without atypical features, neurotic depression, post-traumatic stress disorders and social phobia; dementia of the Alzheimer's type, with early or late onset, with depressed mood; vascular dementia with
15 depressed mood; mood disorders induced by alcohol, amphetamines, cocaine, hallucinogens, inhalants, opioids, phencyclidine, sedatives, hypnotics, anxiolytics and other substances; schizoaffective disorder of the depressed type; and adjustment disorder with depressed mood. Major depressive disorders may also result from a general medical condition including, but not limited to, myocardial infarction, diabetes, miscarriage or
20 abortion, etc.

As used herein, the term anxiety includes anxiety disorders, such as panic disorders with or without agoraphobia, agoraphobia, phobias, for example, social phobias or agoraphobia, obsessive-compulsive disorder, stress disorders including post-traumatic
25 stress disorders, generalised anxiety disorders, acute stress disorders and mixed anxiety-depression disorders

The advantageous profile of anti-anxiety activity obtained by the administration of paroxetine with an NK1 receptor antagonist can be demonstrated in the gerbil social interaction model, according to the method described by Cheeta et al. (Cheeta S. et al.,
30 2001. Brain Research 915: 170-175).

It will be appreciated that the compounds of the combination may be administered simultaneously, either in the same or different pharmaceutical formulations or sequentially. If there is sequential administration, the delay in administering the second
35 and any subsequent active ingredient should not be such as to lose the benefit of a synergistic therapeutic effect of the combination of the active ingredients. It will also be

understood that the compounds of the combination or the physiologically functional derivatives of any thereof, whether presented simultaneously or sequentially, may be administered individually or in multiples or in any combination thereof.

- 5 In a further aspect of the present invention is provided the use of paroxetine and an NK1 receptor antagonist in the manufacture of a medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety, wherein the Nk1 receptor antagonist is selected from
- 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-
10 trifluoromethyl-phenyl)-ethyl]-methyl-amide and physiologically acceptable salts thereof (e.g. methansulphonate and hydrochloride);
- 2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide and physiologically acceptable salts (e.g. hydrochloride, methanesulphonate or maleate);
- 15 2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide and physiologically acceptable salts (e.g. hydrochloride, methanesulphonate or maleate);
- 2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl-benzyl)]-(2S-phenyl-piperidin-3S-yl)-amine).
- 20 In a more preferred embodiment, the present invention provides the use of paroxetine and an NK1 receptor antagonist in the manufacture of a medicament for simultaneous or sequential administration wherein the NK1 receptor antagonist is 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or a physiologically acceptable salt thereof (e.g. methansulphonate or
- 25 hydrochloride).

In a further preferred embodiment, the present invention provides the use of paroxetine and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide or a physiologically acceptable acceptable salt thereof (e.g. methanesulphonate or hydrochloride) in the manufacture of a medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

30

In a further preferred embodiment, the present invention provides the use of paroxetine and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a

35

medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

5 In a further preferred embodiment, the present invention provides the use of paroxetine hydrochloride and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

10 In a further preferred embodiment, the present invention provides the use of paroxetine hydrochloride hemihydrate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

15 In a further preferred embodiment, the present invention provides the use of paroxetine hydrochloride anhydrate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

20 In a further preferred embodiment, the present invention provides the use of paroxetine mesylate and 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methanesulphonate in the manufacture of a medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and/or anxiety.

25 The synergistic effects of the combination of paroxetine and the Nk1 receptor antagonist n may be seen over a wide range of ratios of combination of the components, for example, from 1: 100 to 100: 1 (by weight), such as 1:50 to 50:1 (by weight).

30 Each of the components of the combination of paroxetine and the Nk1 receptor antagonist may thus be employed in a dosage which is therapeutically effective when that component is administered alone. However, in a preferred aspect of the invention, each component may be employed in the combination in a dosage which is less than that

normally expected to produce a fully effective therapeutic response when the component is administered alone.

5 The amount of a combination of paroxetine and NK1 receptor antagonist required to be effective as an antidepressive and/or anti-anxiety may, of course, vary and is ultimately at the discretion of the medical practitioner. The factors to be considered include the route of administration and nature of the formulation, the subject mammal's body weight, age and general condition and the nature and severity of the condition to be treated.

10 In general, a suitable dose of paroxetine for administration to a human for the treatment of depression and/or anxiety as a single therapy may be in a dose of 20 or 400 mg per day, Paroxetine is advantageously administered by oral route once a day.

15 According to the present invention a suitable dose of paroxetine for administration in combination with a Nk1 receptor antagonist to a human for the treatment of depression and/or anxiety may be within the single therapy effective dose range. In a preferred aspect of the invention, the dose of paroxetine administered in combination with an NK1 antagonist may surprisingly be lower than the single therapy dose, in particular from 1 to 20mg per day, preferably from 1 to 10mg per day, and more preferably from 2.5 to 10mg per day..

20 In general, a suitable dose of NK1 receptor antagonist for administration to a human as a single therapy may be in the range of 1 to 300 mg per day, depending upon the particular compound.

25 According to the present invention, a suitable dose of an NK1 receptor antagonist for administration in combination with paroxetine maybe within this singe therapy effective dose range. In a preferred aspect of the invention, the dose of the NK1 receptor antagonist administered in combination with paroxetine may surprisingly be lower than the normal single therapy dose, in particular from 1 to 25 mg per day, preferably from 1 to 10 mg per day and more preferably from 1 to 7.5 mg per day.

30 Combinations according to the invention in which the dosage of the individual paroxetine and NK1 receptor antagonist components are administered below the usual single therapeutic dosages show surprising and synergistic levels of efficacy for the treatment and/or prophylaxis of depression and/or anxiety.

Unless otherwise indicated, all weights of active ingredients are calculated in terms of the drug *per se*. The desired dose may preferably be presented as one, two, three, four, five, six or more sub-doses administered at appropriate intervals throughout the day.

- 5 The components of the combination which may be referred to as active ingredients may be administered for therapy to an animal e.g. a mammal including a human in a conventional manner.

While it is possible for the active ingredients of the combination to be administered as the raw chemical, it is preferable to present them as a pharmaceutical formulation. Pharmaceutical formulations according to the present invention comprise a combination according to the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. The carrier(s) must be acceptable in the sense of being compatible with the other ingredients of the formula and not deleterious to the recipient thereof. When the individual components of the combination are administered separately, they are generally each presented as a pharmaceutical formulation. The references hereinafter to formulations refer, unless otherwise stated, to formulations containing either the combination or a component thereof.

- 20 A combination of paroxetine and a Nk1 receptor antagonist may conveniently be presented as a pharmaceutical formulation in a unitary dosage form.

Pharmaceutical formulations are often prescribed to the patient in "patient packs" containing the whole course of treatment in a single package, usually a blister pack. Patient packs have an advantage over traditional prescriptions, where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve patient compliance with the physician's instructions and, therefore, lead generally to more successful treatment.

It will be understood that the administration of the combination of the invention by means of a single patient pack, or patient packs of each formulation, containing within a package insert instructing the patient to the correct use of the invention is a desirable additional feature of this invention.

According to a further aspect of the invention provided is a multiple, for example, double or triple, pack comprising at least paroxetine and a Nk1 receptor antagonist and an information insert containing directions on the use of the combination of the invention.

- 5 Formulations include those suitable for oral, rectal, nasal, topical (including transdermal, buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. Such methods represent a further feature of the present invention and
- 10 include the step of bringing into association the active ingredients with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product.
- 15 Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, caplets, cachets or tablets each containing a predetermined amount of the active ingredients; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus,
- 20 electuary or paste.

A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as a powder or granules, optionally

25 mixed with a binder (e.g. povidone, gelatin, hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (e.g. sodium starch glycollate, sodium croscarmellose cross-linked povidone, cross-linked sodium carboxymethyl cellulose) surface-active or dispersing agent. Molded tablets may be made by molding a mixture of the powdered compound moistened with an inert liquid diluent in a suitable machine. The tablets may

30 optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredients therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile. Tablets may optionally be provided with an enteric coating, to provide release in parts of the gut other than the stomach.

35

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredients in a flavored base, usually sucrose and acacia or tragacanth;

pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier. Formulations for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or polyethylene glycols.

Topical administration may also be by means of a transdermal iontophoretic device.

Formulations suitable for vaginal administration may be presented as tablets, pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Pharmaceutical formulations suitable for rectal administration wherein the carrier is a solid are most preferably presented as unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be conveniently formed by admixture of the active combination with the softened or melted carrier(s) followed by chilling and shaping in molds.

Formulations suitable for parenteral administration include aqueous and nonaqueous isotonic sterile injection solutions which may contain anti-oxidants, buffers, preservatives and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents; and liposomes or other microparticulate systems which are designed to target the compound to blood components or one or more organs. The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example, those suitable for oral administration may include such further agents as sweeteners, thickeners and flavoring agents.

The pharmaceutical composition of the invention containing the two active ingredients may be prepared according to conventional techniques well known in the pharmaceutical industry. Thus, for example the paroxetine and the NK1 receptor antagonists may be admixed together with suitable excipients such as those described above for the formulation of each of the active ingredients separately. Tablets may be prepared, for example by direct compression of such a mixture or using other conventional methods. Bilayer tablets may be prepared according to conventional procedure. Thus, for example, by separately compressing the two blends in a suitable tableting machine with two filling stations. Capsules may be prepared by filling the blend along with suitable excipients into gelatin capsules, using a suitable filling machine. Controlled release forms for oral or rectal administration may be formulated in a conventional manner associated with controlled release forms.

Biological data:

The advantageous profile of anti anti-anxiety activity obtained by the administration of paroxetine with an NK1 receptor antagonist can be demonstrated in the gerbil social interaction model.

In the experiment 1 the NK1 receptor antagonist used is 2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl-benzyl)]-(2S-phenyl-piperidin-3S-yl)-amine(herein after compound A)

In the experiment 2 the NK1 receptor antagonist used is 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate (herein after compound B)

Experiment 1

Paroxetine (0.3 mg/kg p.o.), compound A (0.03 mg/kg i.p.) and a combination of paroxetine (0.3 mg/kg p.o.) and compound A (0.03 mg/kg i.p.) were administered in mongolian gerbils to assess the effect on time spent in active social interactions.

Experiment 2

Paroxetine (0.3 mg/kg p.o.), compound B (0.1 mg/kg i.p) and a combination of paroxetine (0.3 mg/kg p.o.) and compound B (0.1 mg/kg i.p.) were administered...

The results obtained one hour after administration, express as a percentage variation of the time spent in active social interaction by each animal, in respect to the value obtained by treatment of control animals are summarised below.

Compound			
	paroxetine 0.3mg/kg	Compound A 0.03mg/kg	paroxetine/0.3mg/kg GR205171 0.03mg/kg
% variation	- 2%	+ 19.9%	+ 40.3%

Compound			
	paroxetine 0.3mg/kg	Compound B 0.1mg/kg	paroxetine/0.3mg/kg GW597599 0.1mg/kg
% variation	- 2%	- 8%	+ 56%

The variation of amount of time spent in active social interaction by each animal after treatment of a combination of paroxetine and compound A or paroxetine and compound B is significantly greater than was to be expected from a consideration of the therapeutic response of the components administered separately.

Thus, the above results provide evidence for a synergistic effect between an NK1 receptor antagonist with paroxetine in a social stress assay.

The application of which this description and claims forms part may be used as a basis for priority in respect of any subsequent application. The claims of such subsequent application may be directed to any feature or combination of features described herein. They may take the form of product, composition, process, or use claims and may include, by way of example and without limitation, the following claims:

CLAIMS

1. A composition comprising paroxetine and a Nk1 receptor antagonist.
2. A composition as claimed in claim 1 wherein the Nk1 receptor antagonist is elected from
2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide ;
2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide;
2-(R)-(4-Fluoro-2-methyl-phenyl)-4-(S)-((8aS)-6-oxo-hexahydro-pyrrolo[1,2-a]-pyrazin-2-yl)-piperidine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methanamide
2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl-benzyl)]-(2S-phenyl-piperidin-3S-yl)-amine; and
physiologically acceptable salts thereof.
3. A composition according to claim 1 wherein the Nk1 receptor antagonist is 2-(S)-(4-Fluoro-2-methyl-phenyl)-piperazine-1-carboxylic acid [1-(R)-(3,5-bis-trifluoromethyl-phenyl)-ethyl]-methyl-amide methansulphonate.
4. A composition according to any of claims 1 to 3 for use in the treatment and or prophylaxis of depression and /or anxiety.
5. A method for the treatment and/or prophylaxis of depression and/or anxiety in a mammal including a human, which comprises treating said animal with a therapeutically effective amount of a composition as claimed in any of claims 1 to 3.
6. A method according to any of the claims 1 to 3 wherein the composition is administered as a single combined formulation.
7. A pharmaceutical formulation comprising a composition according to any of the claims 1 to 3 together with one or more pharmaceutically acceptable carriers or excipients.
8. The use of paroxetine and an NK1 receptor antagonist in the manufacture of a medicament for simultaneous or sequential administration for the treatment and/or prophylaxis of depression and /or anxiety.

ABSTRACT

The present invention relates to therapeutic combinations comprising paroxetine and a Nk1 receptor antagonist, to pharmaceutical compositions containing said combinations and their use in the treatment of depression and /or anxiety.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/004122A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/445 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	US 2003/144270 A1 (WROBLESKI MICHELLE LACI ET AL) 31 July 2003 (2003-07-31) page 8, paragraphs 173, 174	1-17
Y	US 2002/123491 A1 (CHEN XIAO ET AL) 5 September 2002 (2002-09-05) page 3, paragraphs 36, 37	1-17
Y	US 2003/064980 A1 (CHEN XIAO ET AL) 3 April 2003 (2003-04-03) page 3, paragraphs 38, 39	1-17
Y	WO 02/10141 A (AHLIJANIAN MICHAEL KIRK ; SANNER MARK ALLEN (US); VILLALOBOS ANABELLA) 7 February 2002 (2002-02-07) page 42, lines 14, 26	1-17
	-/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the International search

19 July 2004

Date of mailing of the International search report

27/07/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Heller, D

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/004122

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01/44200 A (CHEN XIAO ; PALIWAL SUNIL (US); XIAO DONG (US); BLYTHIN DAVID J (US);) 21 June 2001 (2001-06-21) page 5, lines 8-21 -----	1-17
Y	WO 02/32867 A (GLAXO GROUP LTD ; TRANQUILLINI MARIA ELVIRA (IT); MARAGNI PAOLO (IT);) 25 April 2002 (2002-04-25) cited in the application page 6, lines 22,23 page 13, line 8 -----	1-17

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2004/004122

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 15
because they relate to subject matter not required to be searched by this Authority, namely:
Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP2004/004122

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 2003144270	A1	31-07-2003	WO 03042173 A1	22-05-2003
US 2002123491	A1	05-09-2002	US 2003064980 A1	03-04-2003
			AU 2097601 A	25-06-2001
			CA 2393672 A1	21-06-2001
			CN 1434805 T	06-08-2003
			EP 1237874 A2	11-09-2002
			HU 0301607 A2	29-12-2003
			JP 2003522739 T	29-07-2003
			WO 0144200 A2	21-06-2001
US 2003064980	A1	03-04-2003	US 2002123491 A1	05-09-2002
			AU 2097601 A	25-06-2001
			CA 2393672 A1	21-06-2001
			CN 1434805 T	06-08-2003
			EP 1237874 A2	11-09-2002
			HU 0301607 A2	29-12-2003
			JP 2003522739 T	29-07-2003
			WO 0144200 A2	21-06-2001
WO 0210141	A	07-02-2002	AU 7094401 A	13-02-2002
			BG 107469 A	30-09-2003
			BR 0112862 A	01-07-2003
			CA 2418115 A1	07-02-2002
			CN 1444567 T	24-09-2003
			CZ 20030225 A3	18-02-2004
			EP 1305295 A1	02-05-2003
			HR 20030048 A1	30-04-2003
			HU 0303069 A2	01-03-2004
			WO 0210141 A1	07-02-2002
			JP 2004505111 T	19-02-2004
			NO 20030472 A	27-03-2003
			SK 1042003 A3	04-05-2004
			US 2003083352 A1	01-05-2003
			US 2002119963 A1	29-08-2002
WO 0144200	A	21-06-2001	AU 2097601 A	25-06-2001
			CA 2393672 A1	21-06-2001
			CN 1434805 T	06-08-2003
			EP 1237874 A2	11-09-2002
			HU 0301607 A2	29-12-2003
			JP 2003522739 T	29-07-2003
			WO 0144200 A2	21-06-2001
			US 2003064980 A1	03-04-2003
			US 2002123491 A1	05-09-2002
WO 0232867	A	25-04-2002	AU 9572301 A	29-04-2002
			BR 0114637 A	30-09-2003
			CA 2425876 A1	25-04-2002
			CN 1483025 T	17-03-2004
			EP 1326832 A1	16-07-2003
			WO 0232867 A1	25-04-2002
			JP 2004511544 T	15-04-2004
			NO 20031561 A	14-06-2003
			US 2004014770 A1	22-01-2004

This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images
problems checked, please do not report the
problems to the IFW Image Problem Mailbox**